

AFFINE SURFACES WITH TRIVIAL MAKAR-LIMANOV INVARIANT

DANIEL DAIGLE

ABSTRACT. We study the class of 2-dimensional affine \mathbf{k} -domains R satisfying $\text{ML}(R) = \mathbf{k}$, where \mathbf{k} is an arbitrary field of characteristic zero. In particular, we obtain the following result: *Let R be a localization of a polynomial ring in finitely many variables over a field of characteristic zero. If $\text{ML}(R) = K$ for some field $K \subset R$ such that $\text{trdeg}_K R = 2$, then R is K -isomorphic to $K[X, Y, Z]/(XY - P(Z))$ for some nonconstant $P(Z) \in K[Z]$.*

1. INTRODUCTION

Let us recall the definition of the Makar-Limanov invariant:

1.1. Definition. If R is a ring of characteristic zero, a derivation $D : R \rightarrow R$ is said to be *locally nilpotent* if for each $r \in R$ there exists $n \in \mathbb{N}$ (depending on r) such that $D^n(r) = 0$. We use the following notations:

$$\begin{aligned} \text{LND}(R) &= \text{set of locally nilpotent derivations } D : R \rightarrow R \\ \text{KLND}(R) &= \{ \ker D \mid D \in \text{LND}(R) \text{ and } D \neq 0 \} \\ \text{ML}(R) &= \bigcap_{D \in \text{LND}(R)} \ker(D). \end{aligned}$$

We are interested in the class of 2-dimensional affine \mathbf{k} -domains R satisfying $\text{ML}(R) = \mathbf{k}$, where \mathbf{k} is a field of characteristic zero. The corresponding class of affine algebraic surfaces was studied by several authors ([1], [2], [7], [8], [9], [14], [17], in particular), but almost always under the assumption that \mathbf{k} is algebraically closed, or even $\mathbf{k} = \mathbb{C}$. In this paper we obtain some partial results valid when \mathbf{k} is an arbitrary field of characteristic zero. We are particularly interested in the following subclass:

1.2. Definition. Given a field \mathbf{k} of characteristic zero, let $\mathfrak{D}(\mathbf{k})$ be the class of \mathbf{k} -algebras isomorphic to $\mathbf{k}[X, Y, Z]/(XY - \varphi(Z))$ for some nonconstant polynomial in one variable $\varphi(Z) \in \mathbf{k}[Z] \setminus \mathbf{k}$, where X, Y, Z are indeterminates over \mathbf{k} .

The class $\mathfrak{D}(\mathbf{k})$ was studied in [4], [5] and [16], in particular. It is well-known that if $R \in \mathfrak{D}(\mathbf{k})$ then R is a 2-dimensional normal affine domain satisfying $\text{ML}(R) = \mathbf{k}$. It is also known that the converse is not true, which raises the following:

2000 *Mathematics Subject Classification.* Primary: 14R10. Secondary: 14R05, 14R20.

Key words and phrases. Locally nilpotent derivations, group actions, Danielewski surfaces, affine surfaces, Makar-Limanov invariant, absolute constants.

Research supported by a grant from NSERC Canada.

Question. *Suppose that R is a 2-dimensional affine \mathbf{k} -domain with $\text{ML}(R) = \mathbf{k}$. Under what additional assumptions can we infer that $R \in \mathfrak{D}(\mathbf{k})$?*

Section 3 completely answers this question in the case where R is a smooth \mathbf{k} -algebra. This is achieved by reducing to the case $\mathbf{k} = \mathbb{C}$, which was solved by Bandman and Makar-Limanov. This reduction is non-trivial, and makes essential use of the main result of Section 2. Also note Corollary 3.8, which gives a pleasant answer to the above question in the factorial case. Then we derive several consequences from Section 3, for instance consider the following special case of Theorem 4.1:

Let R be a localization of a polynomial ring in finitely many variables over a field of characteristic zero. If $\text{ML}(R) = K$ for some field $K \subset R$ such that $\text{trdeg}_K R = 2$, then $R \in \mathfrak{D}(K)$.

In turn, this has consequences in the study of G_a -actions on \mathbb{C}^n .

Conventions. All rings and algebras are commutative, associative and unital. If A is a ring, we write A^* for the units of A ; if A is a domain, $\text{Frac } A$ is its field of fractions. If $A \subseteq B$ are rings, “ $B = A^{[n]}$ ” means that B is A -isomorphic to the polynomial algebra in n variables over A . If L/K is a field extension, “ $L = K^{(n)}$ ” means that L is a purely transcendental extension of K and $\text{trdeg}_K L = n$ (transcendence degree).

In [5], one defines a Danielewski surface to be a pair (R, \mathbf{k}) such that $R \in \mathfrak{D}(\mathbf{k})$. In the present paper we avoid using the term “Danielewski surface” in that sense, because it is incompatible with accepted usage. The reader should keep this in mind when consulting [5] (our main reference for Section 2).

2. BASE EXTENSION

Let \mathbf{k} be a field of characteristic zero. It is clear that if $R \in \mathfrak{D}(\mathbf{k})$ then $K \otimes_{\mathbf{k}} R \in \mathfrak{D}(K)$ for every field extension K/\mathbf{k} . However, if $K \otimes_{\mathbf{k}} R \in \mathfrak{D}(K)$ for some K , it does not follow that $R \in \mathfrak{D}(\mathbf{k})$ (see Example 2.2, below).

2.1. *Remark.* If $R \in \mathfrak{D}(\mathbf{k})$ then $\text{Spec } R$ has infinitely many \mathbf{k} -rational points. (Indeed, if $R = \mathbf{k}[X, Y, Z]/(XY - \varphi(Z))$ then there is a bijection between the set of \mathbf{k} -rational points of $\text{Spec } R$ and the zero-set in \mathbf{k}^3 of the polynomial $XY - \varphi(Z)$.)

2.2. **Example.** Let $A = \mathbb{R}[X, Y, Z]/(f)$, where $f = X^2 + Y^2 + Z^2$. Viewing f as an element of $\mathbb{C}[X, Y, Z]$ we have $f = (X + iY)(X - iY) + Z^2$ (where $i^2 = -1$), so $\mathbb{C} \otimes_{\mathbb{R}} A \cong \mathbb{C}[U, V, W]/(UV + W^2) \in \mathfrak{D}(\mathbb{C})$. As $\text{Spec } A$ has only one \mathbb{R} -rational point, $A \notin \mathfrak{D}(\mathbb{R})$ by Remark 2.1. Thus

$$A \notin \mathfrak{D}(\mathbb{R}) \text{ and } \mathbb{C} \otimes_{\mathbb{R}} A \in \mathfrak{D}(\mathbb{C}).$$

Note¹ that Theorem 2.3 (below) implies that $\text{ML}(A) = A$. Moreover, if we define $A' = \mathbb{R}[U, V, W]/(UV + W^2) \in \mathfrak{D}(\mathbb{R})$ then $A \not\cong A'$ but $\mathbb{C} \otimes_{\mathbb{R}} A \cong \mathbb{C} \otimes_{\mathbb{R}} A'$.

2.3. **Theorem.** *For an algebra R over a field \mathbf{k} of characteristic zero, the following conditions are equivalent:*

¹A different proof that $\text{ML}(A) = A$ is given in [13, 9.21].

- (a) $R \in \mathfrak{D}(\mathbf{k})$
- (b) $\text{ML}(R) \neq R$ and there exists a field extension K/\mathbf{k} such that $K \otimes_{\mathbf{k}} R \in \mathfrak{D}(K)$.

We shall prove this after some preparation.

2.4. Some facts. Refer to [11] or [13] for background on locally nilpotent derivations. Statement (c) is due to Rentschler [20] and (d) to Nouazé and Gabriel [19] and Wright [21].

- (a) If $A \in \text{KLND}(B)$ where B is a domain of characteristic zero then A is *factorially closed* in B (i.e., if $x, y \in B \setminus \{0\}$ and $xy \in A$ then $x, y \in A$). It follows that $\text{ML}(B)$ is factorially closed in B . Any factorially closed subring A of B is in particular *algebraically closed* in B (i.e., if $x \in B$ is a root of a nonzero polynomial with coefficients in A then $x \in A$) and satisfies $A^* = B^*$ (in particular, any field contained in B is contained in A).
- (b) Let B be a noetherian domain of characteristic zero. If $0 \neq D \in \text{LND}(B)$ then $D = \alpha D_0$ for some $\alpha \in \ker(D)$ and $D_0 \in \text{LND}(B)$ where D_0 is *irreducible* (i.e., the only principal ideal of B which contains $D_0(B)$ is B).
- (c) Let $B = \mathbf{k}^{[2]}$ where \mathbf{k} is a field of characteristic zero. If $D \in \text{LND}(B)$ is irreducible then there exist X, Y such that $B = \mathbf{k}[X, Y]$ and $D = \partial/\partial Y$.
- (d) Let B be a \mathbb{Q} -algebra. If $D \in \text{LND}(B)$ and $s \in B$ satisfy $Ds \in B^*$ then $B = A[s] = A^{[1]}$ where $A = \ker D$.

2.5. Lemma. *Let \mathbf{k} be a field of characteristic zero and R a \mathbf{k} -algebra satisfying:*

there exists a field extension $\bar{\mathbf{k}}/\mathbf{k}$ such that $\bar{\mathbf{k}} \otimes_{\mathbf{k}} R \in \mathfrak{D}(\bar{\mathbf{k}})$.

Then R is a two-dimensional normal affine domain over \mathbf{k} and $R^ = \mathbf{k}^*$.*

Proof. This is rather simple but it will be convenient to refer to this proof later. Choose a field extension $\bar{\mathbf{k}}/\mathbf{k}$ such that $\bar{\mathbf{k}} \otimes_{\mathbf{k}} R \in \mathfrak{D}(\bar{\mathbf{k}})$ and let $\bar{R} = \bar{\mathbf{k}} \otimes_{\mathbf{k}} R$. As R is a flat \mathbf{k} -module, the canonical homomorphism $\mathbf{k} \otimes_{\mathbf{k}} R \rightarrow \bar{\mathbf{k}} \otimes_{\mathbf{k}} R$ is injective, so we may regard R as a subring of \bar{R} . In particular, R is an integral domain and we have the diagram:

$$\begin{array}{ccccccc} \bar{\mathbf{k}} & \hookrightarrow & \bar{R} & \hookrightarrow & S^{-1}\bar{R} & \hookrightarrow & \text{Frac } \bar{R} \\ \uparrow & & \uparrow & & \uparrow & & \\ \mathbf{k} & \hookrightarrow & R & \hookrightarrow & \text{Frac } R & & \end{array}$$

where $S = R \setminus \{0\}$. Let \mathcal{B} be a basis of $\bar{\mathbf{k}}$ over \mathbf{k} such that $1 \in \mathcal{B}$. Note that \mathcal{B} is also a basis of the free R -module \bar{R} and of the vector space $S^{-1}\bar{R}$ over $\text{Frac } R$. It follows:

$$(1) \quad \bar{\mathbf{k}} \cap R = \mathbf{k} \quad \text{and} \quad \bar{R} \cap \text{Frac } R = R.$$

As $\bar{R} \in \mathfrak{D}(\bar{\mathbf{k}})$, [5, 2.3] implies that $\bar{R}^* = \bar{\mathbf{k}}^*$ and that \bar{R} is a normal domain; so (1) implies that $R^* = \mathbf{k}^*$ and that R is a normal domain. Also:

$$(2) \quad \text{If } E \text{ is a subset of } R \text{ such that } \bar{\mathbf{k}}[E] = \bar{R}, \text{ then } \mathbf{k}[E] = R.$$

Indeed, \mathcal{B} is a basis of the R -module \bar{R} and a spanning set of the $\mathbf{k}[E]$ -module \bar{R} ; as $\mathbf{k}[E] \subseteq R$, it follows that $\mathbf{k}[E] = R$.

Note that R is affine over \mathbf{k} , by (2) and the fact that \bar{R} is affine over $\bar{\mathbf{k}}$. Let $n = \dim R$ then, by Noether Normalization Lemma, there exists a subalgebra $R_0 = \mathbf{k}^{[n]}$ of R over which R is integral. Then $\bar{R} = \bar{\mathbf{k}} \otimes_{\mathbf{k}} R$ is integral over $\bar{\mathbf{k}} \otimes_{\mathbf{k}} R_0 = \bar{\mathbf{k}}^{[n]}$, so $n = \dim \bar{R} = 2$. \square

We borrow the following notation from [5, 2.1].

2.6. Definition. Given a \mathbf{k} -algebra R , let $\Gamma_{\mathbf{k}}(R)$ denote the (possibly empty) set of ordered triples $(x_1, x_2, y) \in R \times R \times R$ satisfying:

The \mathbf{k} -homomorphism $\mathbf{k}[X_1, X_2, Y] \rightarrow R$ defined by

$$X_1 \mapsto x_1, X_2 \mapsto x_2 \text{ and } Y \mapsto y$$

is surjective and has kernel equal to $(X_1X_2 - \varphi(Y))\mathbf{k}[X_1, X_2, Y]$ for some nonconstant polynomial in one variable $\varphi(Y) \in \mathbf{k}[Y]$.

Note that $R \in \mathfrak{D}(\mathbf{k})$ if and only if $\Gamma_{\mathbf{k}}(R) \neq \emptyset$.

Proof of Theorem 2.3. That $R \in \mathfrak{D}(\mathbf{k})$ implies $\text{ML}(R) = \mathbf{k}$ is well-known (for instance it follows from part (d) of [5, 2.3]), so it suffices to prove that (b) implies (a).

Suppose that R satisfies (b). Note that if K/\mathbf{k} is a field extension satisfying $K \otimes_{\mathbf{k}} R \in \mathfrak{D}(K)$ then for any field extension L/K we have $L \otimes_{\mathbf{k}} R \in \mathfrak{D}(L)$. In particular, there exists a field extension $\bar{\mathbf{k}}/\mathbf{k}$ such that $\bar{\mathbf{k}} \otimes_{\mathbf{k}} R \in \mathfrak{D}(\bar{\mathbf{k}})$ and such that $\bar{\mathbf{k}}$ is an algebraically closed field. We fix such a field $\bar{\mathbf{k}}$. The fact that $\bar{\mathbf{k}}$ is algebraically closed implies that

$$(3) \quad \text{the fixed field } \bar{\mathbf{k}}^G \text{ is equal to } \mathbf{k}$$

where $G = \text{Gal}(\bar{\mathbf{k}}/\mathbf{k})$. We use the notation $(\bar{R}, \mathcal{B}, \text{etc})$ introduced in the proof of Lemma 2.5. As $\text{ML}(R) \neq R$, there exists $0 \neq D \in \text{LND}(R)$. Let $\bar{D} \in \text{LND}(\bar{R})$ be the unique extension of D , let $A = \ker D$ and $\bar{A} = \ker \bar{D}$.

It follows from [5] that $\bar{A} = \bar{\mathbf{k}}^{[1]}$ ([5, 2.3] shows that some element of $\text{KLND}(\bar{R})$ is a $\bar{\mathbf{k}}^{[1]}$ and, by [5, 2.7.2], $\text{Aut}_{\bar{\mathbf{k}}}(\bar{R})$ acts transitively on $\text{KLND}(\bar{R})$). Applying the exact functor $\bar{\mathbf{k}} \otimes_{\mathbf{k}} _$ to the exact sequence $0 \rightarrow A \rightarrow R \xrightarrow{D} R$ of \mathbf{k} -linear maps shows that $\bar{\mathbf{k}} \otimes_{\mathbf{k}} A = \bar{A} = \bar{\mathbf{k}}^{[1]}$, so $A = \mathbf{k}^{[1]}$. Choose $f \in R$ such that $A = \mathbf{k}[f]$, then $\bar{A} = \bar{\mathbf{k}}[f]$.

Consider the nonzero ideals $I = A \cap D(R)$ and $\bar{I} = \bar{A} \cap \bar{D}(\bar{R})$ of A and \bar{A} respectively. Let $\psi \in A$ and $s \in R$ be such that $I = \psi A$ and $D(s) = \psi$. We claim that

$$(4) \quad \bar{I} = \psi \bar{A}.$$

Indeed, an arbitrary element of \bar{I} is of the form $\bar{D}(\sigma)$ where $\sigma \in \bar{R}$ and $\bar{D}^2(\sigma) = 0$. Write $\sigma = \sum_{\lambda \in \mathcal{B}} s_{\lambda} \lambda$ with $s_{\lambda} \in R$, then $0 = \bar{D}^2(\sigma) = \sum_{\lambda \in \mathcal{B}} D^2(s_{\lambda}) \lambda$, so for all $\lambda \in \mathcal{B}$ we have $D^2(s_{\lambda}) = 0$, hence $D(s_{\lambda}) \in I = \psi A$, and consequently $\bar{D}(\sigma) \in \psi \bar{A}$, which proves (4).

By 2.4(b), $\bar{D} = \alpha \Delta$ for some $\alpha \in \bar{A} \setminus \{0\}$ and some irreducible $\Delta \in \text{LND}(\bar{R})$. Consider the nonzero ideal $I_0 = \bar{A} \cap \Delta(\bar{R})$ of \bar{A} . We claim that

$$(5) \quad I_0 = \Delta(s) \bar{A}.$$

To see this, consider an arbitrary element $\Delta(\sigma)$ of I_0 (where $\sigma \in \bar{R}$, $\Delta^2(\sigma) = 0$). Then $\alpha\Delta(\sigma) = \bar{D}(\sigma) \in \bar{I} = \psi\bar{A} = \bar{D}(s)\bar{A} = \alpha\Delta(s)\bar{A}$, so $\Delta(\sigma) \in \Delta(s)\bar{A}$ and (5) is proved.

Consider the case where $\Delta(s) \in \bar{R}^*$. Then $\bar{R} = \bar{A}[s] = \bar{\mathbf{k}}[f, s]$ by 2.4(d), so (2) implies that $R = \mathbf{k}[f, s] = \mathbf{k}^{[2]}$, so in particular $R \in \mathfrak{D}(\mathbf{k})$ and we are done.

From now-on assume that $\Delta(s) \notin \bar{R}^*$. By [5, 2.8], $\bar{A} = \bar{\mathbf{k}}[\Delta(y)]$ for some $y \in \bar{R}$. Note that $\Delta(y) \in I_0$, so (5) gives $\Delta(s) \mid \Delta(y)$ in \bar{A} . As $\Delta(y)$ is an irreducible element of \bar{A} (because $\bar{\mathbf{k}}[\Delta(y)] = \bar{A} = \bar{\mathbf{k}}^{[1]}$) and $\Delta(s) \notin \bar{A}^*$, we have $\bar{\mathbf{k}}[\Delta(s)] = \bar{A} = \bar{\mathbf{k}}[f]$ and consequently $\Delta(s) = \mu(f - \lambda)$ for some $\mu \in \bar{\mathbf{k}}^*$, $\lambda \in \bar{\mathbf{k}}$. We may as well replace Δ by $\mu^{-1}\Delta$, so

$$(6) \quad \Delta(s) = f - \lambda, \quad \text{for some } \lambda \in \bar{\mathbf{k}}.$$

We claim:

$$(7) \quad \{c \in \bar{\mathbf{k}} \mid \bar{R}/(f - c)\bar{R} \text{ is not an integral domain}\} = \{\lambda\}.$$

Indeed, [5, 2.8] implies that there exists $x_2 \in \bar{R}$ such that $(f - \lambda, x_2, s) \in \Gamma_{\bar{\mathbf{k}}}(\bar{R})$. This means (cf. 2.6) that the $\bar{\mathbf{k}}$ -homomorphism $\pi : \bar{\mathbf{k}}[X_1, X_2, Y] \rightarrow \bar{R}$ defined by $X_1 \mapsto f - \lambda$, $X_2 \mapsto x_2$, $Y \mapsto s$, is surjective and has kernel $(X_1X_2 - P(Y))$ for some nonconstant $P(Y) \in \bar{\mathbf{k}}[Y]$ (where X_1, X_2, Y are indeterminates). By (5) and $\Delta(s) \notin \bar{R}^*$, we see that there does not exist $\sigma \in \bar{R}$ such that $\Delta(\sigma) = 1$; as Δ is irreducible, it follows from 2.4(c) that $\bar{R} \neq \bar{\mathbf{k}}^{[2]}$ and hence that $\deg_Y P(Y) > 1$. Thus, for $c \in \bar{\mathbf{k}}$,

$$\bar{R}/(f - c)\bar{R} \cong \bar{\mathbf{k}}[X_1, X_2, Y]/(X_1 - (c - \lambda), X_1X_2 - P(Y))$$

is a domain if and only if $c \neq \lambda$. This proves (7).

Let $\theta \in \text{Gal}(\bar{\mathbf{k}}/\mathbf{k})$. Then θ extends to some $\Theta \in \text{Aut}_R(\bar{R})$ and Θ determines a ring isomorphism

$$\bar{R}/(f - \lambda)\bar{R} \cong \bar{R}/\Theta(f - \lambda)\bar{R} = \bar{R}/(f - \theta(\lambda))\bar{R}.$$

So $\bar{R}/(f - \theta(\lambda))\bar{R}$ is not a domain and it follows from (7) that $\theta(\lambda) = \lambda$. As this holds for every $\theta \in \text{Gal}(\bar{\mathbf{k}}/\mathbf{k})$, (3) implies that $\lambda \in \mathbf{k}$. To summarize, if we define $x_1 = f - \lambda$ then

$$x_1, s \in R \text{ and there exists } x_2 \in \bar{R} \text{ such that } (x_1, x_2, s) \in \Gamma_{\bar{\mathbf{k}}}(\bar{R}).$$

We now show that x_2 can be chosen in R . Consider the ideals $J = \mathbf{k}[s] \cap x_1R$ of $\mathbf{k}[s]$ and $\bar{J} = \bar{\mathbf{k}}[s] \cap x_1\bar{R}$ of $\bar{\mathbf{k}}[s]$, and choose $\varphi(Y) \in \mathbf{k}[Y]$ such that $J = \varphi(s)\mathbf{k}[s]$. Let $\Phi(s)$ be any element of \bar{J} (where $\Phi(Y) \in \bar{\mathbf{k}}[Y]$). Then $\Phi(s) = x_1G$ for some $G \in \bar{R}$. As \mathcal{B} is a basis of the R -module \bar{R} and also of the $\mathbf{k}[Y]$ -module $\bar{\mathbf{k}}[Y]$, we may write $G = \sum_{\lambda \in \mathcal{B}} G_\lambda \lambda$ (where $G_\lambda \in R$) and $\Phi = \sum_{\lambda \in \mathcal{B}} \Phi_\lambda \lambda$ (where $\Phi_\lambda \in \bar{\mathbf{k}}[Y]$). Then $\sum_{\lambda \in \mathcal{B}} (x_1G_\lambda)\lambda = \Phi(s) = \sum_{\lambda \in \mathcal{B}} \Phi_\lambda(s)\lambda$, so for every $\lambda \in \mathcal{B}$ we have $\Phi_\lambda(s) = x_1G_\lambda$, i.e., $\Phi_\lambda(s) \in J = \varphi(s)\mathbf{k}[s]$. We obtain that $\Phi(s) \in \varphi(s)\bar{\mathbf{k}}[s]$, so:

$$\bar{J} = \varphi(s)\bar{\mathbf{k}}[s].$$

On the other hand, [5, 2.4] asserts that $\bar{J} = x_1x_2\bar{\mathbf{k}}[s]$, so $x_1x_2 = \mu\varphi(s)$ for some $\mu \in \bar{\mathbf{k}}^*$. It is clear that if (x_1, x_2, s) belongs to $\Gamma_{\bar{\mathbf{k}}}(\bar{R})$ then so does $(x_1, \mu^{-1}x_2, s)$; so there exists

$x_2 \in \bar{R}$ such that $(x_1, x_2, s) \in \Gamma_{\bar{\mathbf{k}}}(\bar{R})$ and $x_1 x_2 = \varphi(s)$. As $x_2 = \varphi(s)/x_1 \in \text{Frac } R$, (1) implies that $x_2 \in R$. Thus

$$(x_1, x_2, s) \in \Gamma_{\bar{\mathbf{k}}}(\bar{R}), \text{ where } x_1, x_2, s \in R.$$

In particular we have $\bar{R} = \bar{\mathbf{k}}[x_1, x_2, s]$, so (2) gives $R = \mathbf{k}[x_1, x_2, s]$. As $x_1 x_2 = \varphi(s)$ where $\varphi(Y) \in \mathbf{k}[Y]$ is nonconstant, it follows that $(x_1, x_2, s) \in \Gamma_{\mathbf{k}}(R)$ and hence that $R \in \mathfrak{D}(\mathbf{k})$. \square

3. ON A RESULT OF BANDMAN AND MAKAR-LIMANOV

In this paper we adopt the following:

3.1. Definition. Let R be an affine algebra over a field \mathbf{k} and let $q = \dim R$. We say that R is a *complete intersection over \mathbf{k}* if $R \cong \mathbf{k}[X_1, \dots, X_{p+q}]/(f_1, \dots, f_p)$ for some $p \geq 0$ and some $f_1, \dots, f_p \in \mathbf{k}[X_1, \dots, X_{p+q}]$.


We refer to [18, 28.D] for the definition of a *smooth \mathbf{k} -algebra* and to [18, 26.C] for the definition of the R -module $\Omega_{R/\mathbf{k}}$ (the module of differentials of R over \mathbf{k}), where R is a \mathbf{k} -algebra.

3.2. Theorem. *Let \mathbf{k} be a field of characteristic zero and R a smooth affine \mathbf{k} -domain of dimension 2 such that $\text{ML}(R) = \mathbf{k}$. Then the following are equivalent:*

- (a) $R \in \mathfrak{D}(\mathbf{k})$
- (b) R is generated by 3 elements as a \mathbf{k} -algebra
- (c) R is a complete intersection over \mathbf{k}
- (d) $\bigwedge^2 \Omega_{R/\mathbf{k}} \cong R$.

We shall prove this by reducing to the case $\mathbf{k} = \mathbb{C}$, which was proved by Bandman and Makar-Limanov in [1]. That reduction makes essential use of Theorem 2.3.

3.3. Remark. Let \mathbf{k} be a field of characteristic zero. According to the definition of ‘‘Danielewski surface over \mathbf{k} ’’ given in [10], one has the following situation:



where $\text{DANML}(\mathbf{k})$ is the class of Danielewski surfaces S over \mathbf{k} satisfying $\text{ML}(S) = \mathbf{k}$, $\text{SML}(\mathbf{k})$ is the larger class of smooth affine surfaces S over \mathbf{k} satisfying $\text{ML}(S) = \mathbf{k}$, and $\mathfrak{D}(\mathbf{k})$ is the class of surfaces corresponding to the already defined class $\mathfrak{D}(\mathbf{k})$ of \mathbf{k} -algebras. Among other things, paper [10] classifies the elements of $\text{DANML}(\mathbf{k})$ and characterizes those which belong to $\mathfrak{D}(\mathbf{k})$. In contrast, Theorem 3.2 characterizes the elements of $\text{SML}(\mathbf{k})$ which belong to $\mathfrak{D}(\mathbf{k})$.

3.4. *Remark.* Let R be a q -dimensional smooth affine domain over a field \mathbf{k} of characteristic zero. Then $X = \text{Spec } R$ is in particular an irreducible regular scheme of finite type over the perfect field \mathbf{k} ; so, by [15, ex. 8.1(c), p. 187], the sheaf of differentials $\Omega_{X/\mathbf{k}}$ is locally free of rank q ; so the canonical sheaf $\omega_X = \bigwedge^q \Omega_{X/\mathbf{k}}$ is locally free of rank 1, i.e., is an invertible sheaf on X . As ω_X and the structure sheaf \mathcal{O}_X are respectively the sheaves associated to the R -modules $\bigwedge^q \Omega_{R/\mathbf{k}}$ and R , the condition $\bigwedge^q \Omega_{R/\mathbf{k}} \cong R$ is equivalent to $\omega_X \cong \mathcal{O}_X$ (one says that X has trivial canonical sheaf). This is also equivalent to the canonical divisor of X being linearly equivalent to zero (because $\text{Pic}(X) \cong \text{Cl}(X)$ by [15, 6.16 p. 145]).

3.5. *Remark.* Let A' and B be algebras over a ring A and let $B' = A' \otimes_A B$. Then $\Omega_{B'/A'} \cong B' \otimes_B \Omega_{B/A}$ (cf. [18, p. 186]) and, for any B -module M , $\bigwedge^n (B' \otimes_B M) \cong B' \otimes_B \bigwedge^n M$ for every n ([3], Chap. 3, §7, No 5, Prop. 8). Consequently, $\bigwedge^n \Omega_{B'/A'} \cong B' \otimes_B \bigwedge^n \Omega_{B/A}$.

3.6. **Lemma.** *Let R be an algebra over a field \mathbf{k} . If R is a complete intersection over \mathbf{k} and a smooth \mathbf{k} -algebra, then $\bigwedge^q \Omega_{R/\mathbf{k}} \cong R$ where $q = \dim R$.*

This is the well-known fact that a smooth complete intersection has trivial canonical sheaf, but we don't know a suitable reference so we sketch a proof.

Proof of 3.6. Let $R = \mathbf{k}[X_1, \dots, X_{p+q}]/(f_1, \dots, f_p)$ and let $\varphi_{ij} \in R$ be the image of $\frac{\partial f_j}{\partial X_i}$. Because R is smooth over \mathbf{k} , [18, 29.E] implies that the matrix (φ_{ij}) satisfies:

$$(8) \quad \text{the } p \times p \text{ determinants of } (\varphi_{ij}) \text{ generate the unit ideal of } R.$$

By [15, 8.4A, p. 173], there is an exact sequence $R^p \xrightarrow{\varphi} R^{p+q} \rightarrow \Omega_{R/\mathbf{k}} \rightarrow 0$ of R -linear maps where φ is the map corresponding to the matrix (φ_{ij}) . Now if R is a ring and $R^p \xrightarrow{\varphi} R^{p+q} \rightarrow M \rightarrow 0$ is an exact sequence of R -linear maps such that φ satisfies (8), then $\bigwedge^q M \cong R$. \square

3.7. **Lemma.** *Let R be an integral domain containing a field \mathbf{k} of characteristic zero. If R is normal and $\text{ML}(R) = \mathbf{k}$, then for any field extension K of \mathbf{k} we have:*

- (a) $K \otimes_{\mathbf{k}} R$ is an integral domain
- (b) $\text{ML}(K \otimes_{\mathbf{k}} R) = K$.

Proof. As $\mathbf{k} = \text{ML}(R)$ is algebraically closed in R (2.4(a)) and R is normal, it follows that \mathbf{k} is algebraically closed in $L = \text{Frac } R$. By [22, Cor. 2, p. 198], $K \otimes_{\mathbf{k}} L$ is an integral domain. As K is flat over \mathbf{k} and $R \rightarrow L$ is injective, $K \otimes_{\mathbf{k}} R \rightarrow K \otimes_{\mathbf{k}} L$ is injective and (a) is proved.

Let $\xi \in \text{ML}(K \otimes_{\mathbf{k}} R)$. Consider a basis \mathcal{B} of K over \mathbf{k} ; note that \mathcal{B} is also a basis of the free R -module $R' = K \otimes_{\mathbf{k}} R$ and write $\xi = \sum_{\lambda \in \mathcal{B}} x_{\lambda} \lambda$ (where $x_{\lambda} \in R$). If $D \in \text{LND}(R)$ then D extends to an element $D' \in \text{LND}(R')$ and the equation $0 = D'(\xi) = \sum_{\lambda \in \mathcal{B}} D(x_{\lambda}) \lambda$ shows that $D(x_{\lambda}) = 0$ for all $\lambda \in \mathcal{B}$. As this holds for every $D \in \text{LND}(R)$, we have $x_{\lambda} \in \text{ML}(R) = \mathbf{k}$ for all λ , so $\xi \in K$. \square

Proof of Theorem 3.2. Implications (a) \Rightarrow (b) \Rightarrow (c) are trivial and (c) \Rightarrow (d) is Lemma 3.6, so only (d) \Rightarrow (a) requires a proof. Assume for a moment that $\mathbf{k} = \mathbb{C}$ and suppose that R satisfies (d). Then Lemmas 4 and 5 of [1] imply that $R \in \mathfrak{D}(\mathbb{C})$, so the Theorem is valid in the case $\mathbf{k} = \mathbb{C}$.

Let \mathbf{k} be a field of characteristic zero, consider a smooth affine \mathbf{k} -domain R of dimension 2 such that $\text{ML}(R) = \mathbf{k}$, and suppose that R satisfies (d).

We have $R \cong \mathbf{k}[X_1, \dots, X_n]/(f_1, \dots, f_m)$ for some $m, n \geq 0$ and some $f_1, \dots, f_m \in \mathbf{k}[X_1, \dots, X_n]$. Also consider $D_1, D_2 \in \text{LND}(R)$ such that $\ker D_1 \cap \ker D_2 = \mathbf{k}$. Each D_i can be lifted to a (not necessarily locally nilpotent) \mathbf{k} -derivation δ_i of $\mathbf{k}[X_1, \dots, X_n]$. Let \mathbf{k}_0 be a subfield of \mathbf{k} which is finitely generated over \mathbb{Q} and which contains all coefficients of the polynomials f_i and $\delta_i(X_j)$. Define $R_0 = \mathbf{k}_0[X_1, \dots, X_n]/(f_1, \dots, f_m)$ and note that $\mathbf{k} \otimes_{\mathbf{k}_0} R_0 \cong R$. As $\mathbf{k}_0 \rightarrow \mathbf{k}$ is injective and R_0 is flat over \mathbf{k}_0 , $\mathbf{k}_0 \otimes_{\mathbf{k}_0} R_0 \rightarrow \mathbf{k} \otimes_{\mathbf{k}_0} R_0$ is injective and we may regard R_0 as a subring of R . In particular, R_0 is a domain (a 2-dimensional affine \mathbf{k}_0 -domain). Also note that $D_i(R_0) \subseteq R_0$ for $i = 1, 2$; if $d_i : R_0 \rightarrow R_0$ is the restriction of D_i then $d_1, d_2 \in \text{LND}(R_0)$ and $\ker d_1 \cap \ker d_2 = \mathbf{k} \cap R_0 = \mathbf{k}_0$ (see (1) for the last equality), showing that $\text{ML}(R_0) = \mathbf{k}_0$. As \mathbf{k}_0 is a field and $\mathbf{k} \rightarrow R$ is obtained from $\mathbf{k}_0 \rightarrow R_0$ by base extension, the fact that $\mathbf{k} \rightarrow R$ is smooth implies that $\mathbf{k}_0 \rightarrow R_0$ is smooth (cf. [18, 28.O]).

Consider the R -module $M = \bigwedge^2 \Omega_{R/\mathbf{k}}$ and the R_0 -module $M_0 = \bigwedge^2 \Omega_{R_0/\mathbf{k}_0}$. Consider an isomorphism of R -modules $\theta : R \rightarrow M$ and let $\omega = \theta(1)$. We have $R \otimes_{R_0} M_0 \cong M$ by 3.5, so there is a natural homomorphism $M_0 \rightarrow R \otimes_{R_0} M_0 \cong M$, $x \mapsto 1 \otimes x$; by adjoining a finite subset of \mathbf{k} to \mathbf{k}_0 , we may arrange that there exists $\omega_0 \in M_0$ such that $1 \otimes \omega_0 = \omega$. Consider the R_0 -linear map $f : R_0 \rightarrow M_0$, $f(a) = a\omega_0$. Note that $R = \mathbf{k} \otimes_{\mathbf{k}_0} R_0$ is faithfully flat as an R_0 -module and that applying the functor $R \otimes_{R_0} _$ to f yields the isomorphism θ ; so f is an isomorphism, so $\bigwedge^2 \Omega_{R_0/\mathbf{k}_0} \cong R_0$. As $R \in \mathfrak{D}(\mathbf{k})$ would follow from $R_0 \in \mathfrak{D}(\mathbf{k}_0)$, the problem reduces to proving the case $\mathbf{k} = \mathbf{k}_0$ of the theorem. Now \mathbf{k}_0 is isomorphic to a subfield of \mathbb{C} , so it suffices to prove the theorem in the case $\mathbf{k} \subseteq \mathbb{C}$.

Assume that $\mathbf{k} \subseteq \mathbb{C}$. As R is smooth over \mathbf{k} , the local ring $R_{\mathfrak{p}}$ is regular for every $\mathfrak{p} \in \text{Spec } R$ (by [18, 28.E,F,K]) so in particular R is a normal domain. Then it follows from 3.7 that $R' = \mathbb{C} \otimes_{\mathbf{k}} R$ is an integral domain and that $\text{ML}(R') = \mathbb{C}$. By [18, 28.G], R' is smooth over \mathbb{C} . It is clear that $\dim R' = 2$ (for instance see the proof of 2.5) and 3.5 gives $\bigwedge^2 \Omega_{R'/\mathbb{C}} \cong R' \otimes_R \bigwedge^2 \Omega_{R/\mathbf{k}} \cong R' \otimes_R R \cong R'$. As the Theorem is valid over \mathbb{C} , it follows that $R' \in \mathfrak{D}(\mathbb{C})$. As $\text{ML}(R) = \mathbf{k} \neq R$, Theorem 2.3 implies that $R \in \mathfrak{D}(\mathbf{k})$. \square

3.8. Corollary. *Let R be a 2-dimensional affine domain over a field \mathbf{k} of characteristic zero. If R is a UFD and a smooth \mathbf{k} -algebra satisfying $\text{ML}(R) = \mathbf{k}$, then $R \in \mathfrak{D}(\mathbf{k})$.*

Proof. Since R is a UFD, the scheme $X = \text{Spec } R$ has a trivial divisor class group [15, 6.2 p. 131]. By Remark 3.4, it follows that $\bigwedge^2 \Omega_{R/\mathbf{k}} \cong R$ and the desired conclusion follows from Theorem 3.2. \square

4. LOCALIZATIONS OF NICE RINGS

Throughout this section we fix a field \mathbf{k} of characteristic zero and we consider the class $\mathcal{N}(\mathbf{k})$ of \mathbf{k} -algebras B satisfying the following conditions:

B is a geometrically integral affine \mathbf{k} -domain which is smooth over \mathbf{k} and satisfies at least one of the following conditions:

- *B is a UFD; or*
- *B is a complete intersection over \mathbf{k} .*

Note that $\mathbf{k}^{[n]} \in \mathcal{N}(\mathbf{k})$ for every n .

4.1. Theorem. *Suppose that R is a localization of a ring belonging to the class $\mathcal{N}(\mathbf{k})$. If $\text{ML}(R) = K$ for some field $K \subset R$ such that $\text{trdeg}_K R = 2$, then $R \in \mathfrak{D}(K)$.*

4.2. Lemma. *Let $B \in \mathcal{N}(\mathbf{k})$, let E be a finitely generated \mathbf{k} -subalgebra of B and let $S = E \setminus \{0\}$. Then $S^{-1}B$ is a smooth algebra over the field $S^{-1}E$.*

Proof. Let $\bar{\mathbf{k}}$ be an algebraic closure of \mathbf{k} and define $\bar{E} = \bar{\mathbf{k}} \otimes_{\mathbf{k}} E$ and $\bar{B} = \bar{\mathbf{k}} \otimes_{\mathbf{k}} B$. Note that \bar{B} is a domain because B is geometrically integral, and $\bar{E} \rightarrow \bar{B}$ is injective because $\bar{\mathbf{k}}$ is flat over \mathbf{k} . Let $K = \text{Frac } E$ and $L = \text{Frac } \bar{E}$. As \bar{B} is smooth over $\bar{\mathbf{k}}$, applying [15, 10.7, p. 272] to $\text{Spec } \bar{B} \rightarrow \text{Spec } \bar{E}$ implies that $L \rightarrow L \otimes_{\bar{E}} \bar{B}$ is smooth. It is not difficult to see that $L \rightarrow L \otimes_{\bar{E}} \bar{B}$ is obtained from $K \rightarrow K \otimes_E B$ by base extension. As K is a field and $L \rightarrow L \otimes_{\bar{E}} \bar{B}$ is smooth, it follows from [18, 28.O] that $K \rightarrow K \otimes_E B$ is smooth. \square

4.3. Lemma. *Let $B \in \mathcal{N}(\mathbf{k})$, let S be a multiplicative subset of B and suppose that K is a field such that $\mathbf{k} \cup S \subseteq K \subseteq S^{-1}B$. Then $S^{-1}B$ is a smooth K -algebra and some transcendence basis of K/\mathbf{k} is a subset of B .*

Proof. Note that K/\mathbf{k} is a finitely generated field extension and write $K = \mathbf{k}(\alpha_1, \dots, \alpha_m)$. For each i we have $\alpha_i = b_i/s_i$ for some $b_i \in B$ and $s_i \in S$; as $S \subseteq K$, we have $b_i = s_i \alpha_i \in K$. Define $E = \mathbf{k}[b_1, \dots, b_m, s_1, \dots, s_m] \subseteq K$ and $S_1 = E \setminus \{0\}$, then $S_1^{-1}E = K$ and hence $S_1^{-1}B = S^{-1}B$. By Lemma 4.2, $S^{-1}B$ is a smooth K -algebra. Moreover, $\{b_1, \dots, b_m, s_1, \dots, s_m\}$ contains a transcendence basis of K/\mathbf{k} . \square

Proof of Theorem 4.1. We have $R = S^{-1}B$ for some $B \in \mathcal{N}(\mathbf{k})$ and some multiplicative subset S of B . As $\mathbf{k}^* \cup S \subseteq R^* \subseteq \text{ML}(R) = K$, R is smooth over K by Lemma 4.3. By definition of $\mathcal{N}(\mathbf{k})$, B is a UFD or a complete intersection over \mathbf{k} .

If B is a UFD then so is R ; in this case we obtain $R \in \mathfrak{D}(K)$ by Corollary 3.8, so we are done.

From now-on, assume that B is a complete intersection over \mathbf{k} . Let $q = \dim B$ and write $B = \mathbf{k}[X_1, \dots, X_{p+q}]/(G_1, \dots, G_p)$. Using Lemma 4.3 again, choose a transcendence basis $\{f_1, \dots, f_{q-2}\}$ of K over \mathbf{k} such that $f_1, \dots, f_{q-2} \in B$; let $S_0 = \mathbf{k}[f_1, \dots, f_{q-2}] \setminus \{0\}$ and $K_0 = \mathbf{k}(f_1, \dots, f_{q-2})$. We claim:

$$(9) \quad S_0^{-1}B \text{ is a complete intersection over } K_0.$$

Let us prove this. For $1 \leq i \leq q-2$, choose $F_i \in \mathbf{k}[X_1, \dots, X_{p+q}]$ such that $\pi(F_i) = f_i$ where $\pi : \mathbf{k}[X_1, \dots, X_{p+q}] \rightarrow B$ is the canonical epimorphism. Also, let T_1, \dots, T_{q-2} be extra indeterminates. The \mathbf{k} -homomorphism $\mathbf{k}[T_1, \dots, T_{q-2}, X_1, \dots, X_{p+q}] \rightarrow B$ which maps T_i to f_i and X_i to $\pi(X_i)$ has kernel $(G_1, \dots, G_p, F_1 - T_1, \dots, F_{q-2} - T_{q-2})$, so there is an isomorphism of \mathbf{k} -algebras

$$B \cong \mathbf{k}[T_1, \dots, T_{q-2}, X_1, \dots, X_{p+q}]/(G_1, \dots, G_p, F_1 - T_1, \dots, F_{q-2} - T_{q-2}).$$

Localization gives an isomorphism of \mathbf{k} -algebras

$$(10) \quad S_0^{-1}B \cong \mathbf{k}(T_1, \dots, T_{q-2})[X_1, \dots, X_{p+q}]/(G_1, \dots, G_p, F_1 - T_1, \dots, F_{q-2} - T_{q-2})$$

which maps K_0 onto $\mathbf{k}(T_1, \dots, T_{q-2})$. As the right hand side of (10) is a complete intersection over $\mathbf{k}(T_1, \dots, T_{q-2})$, assertion (9) is proved. Then we obtain

$$(11) \quad \bigwedge^2 \Omega_{S_0^{-1}B/K_0} \cong S_0^{-1}B$$

by Lemma 3.6, because $S_0^{-1}B$ is a smooth K_0 -algebra by Lemma 4.2.

Each element of K belongs to $\text{Frac}(S_0^{-1}B)$ and is algebraic over K_0 , hence integral over $S_0^{-1}B$; as $S_0^{-1}B$ is normal, $K \subseteq S_0^{-1}B$ and hence $S_0^{-1}B = R$. We may therefore rewrite (11) as:

$$(12) \quad \bigwedge^2 \Omega_{R/K_0} \cong R.$$

Applying [18, 26.H] to $K_0 \subseteq K \subseteq R$ gives the exact sequence of R -modules

$$\Omega_{K/K_0} \otimes_K R \rightarrow \Omega_{R/K_0} \rightarrow \Omega_{R/K} \rightarrow 0,$$

where $\Omega_{K/K_0} = 0$ by [18, 27.B]. So $\Omega_{R/K} \cong \Omega_{R/K_0}$ and hence (12) gives $\bigwedge^2 \Omega_{R/K} \cong R$. So $R \in \mathfrak{D}(K)$ by Theorem 3.2. \square

Let \mathbf{k} be a field of characteristic zero, let $B \in \mathcal{N}(\mathbf{k})$ and consider locally nilpotent derivations $D : B \rightarrow B$. See 1.1 for the definition of $\text{KLND}(B)$. It is known that if $A \in \text{KLND}(B)$ then $\text{trdeg}_A(B) = 1$, and if A_1, A_2 are distinct elements of $\text{KLND}(B)$ then $\text{trdeg}_{A_1 \cap A_2}(B) \geq 2$. We are interested in the situation where $\text{trdeg}_{A_1 \cap A_2}(B) = 2$, i.e., when A_1, A_2 are distinct and have an intersection which is as large as possible.

4.4. Corollary. *Let $B \in \mathcal{N}(\mathbf{k})$, where \mathbf{k} is a field of characteristic zero. If $A_1, A_2 \in \text{KLND}(B)$ are such that $\text{trdeg}_{A_1 \cap A_2}(B) = 2$, then the following hold.*

- (a) *Let $R = A_1 \cap A_2$ and $K = \text{Frac } R$. Then $K \otimes_R B \in \mathfrak{D}(K)$.*
- (b) *If B is a UFD then there exists a finite sequence of local slice constructions which transforms A_1 into A_2 .*

Remark. This generalizes results 1.10 and 1.13 of [6]. Local slice construction was originally defined in [12] in the case $B = \mathbf{k}^{[3]}$, and was later generalized in [5].

Proof of Corollary 4.4. Let $S = R \setminus \{0\}$, $\mathcal{A}_i = S^{-1}A_i$ ($i = 1, 2$) and $\mathcal{B} = S^{-1}B = K \otimes_R B$. If $D_i \in \text{LND}(B)$ has kernel A_i , then $S^{-1}D_i \in \text{LND}(\mathcal{B})$ has kernel \mathcal{A}_i ; thus $\mathcal{A}_1, \mathcal{A}_2 \in \text{KLND}(\mathcal{B})$. Using that A_1, A_2 are factorially closed in B , we obtain $\mathcal{A}_1 \cap \mathcal{A}_2 \subseteq K$, so $\text{ML}(\mathcal{B}) \subseteq K$. The reverse inclusion is trivial ($K^* \subseteq \mathcal{B}^* \subseteq \text{ML}(\mathcal{B})$), so $\text{ML}(\mathcal{B}) = K$. Then $\mathcal{B} \in \mathfrak{D}(K)$ by Theorem 4.1, so assertion (a) is proved.

In [5, 3.3], one defines a graph $\underline{\text{KLND}}(B)$ whose vertex-set is $\text{KLND}(B)$; then, given $A, A' \in \text{KLND}(B)$, one says that A' can be obtained from A “by a local slice construction” if there exists an edge in $\underline{\text{KLND}}(B)$ joining vertices A and A' . So assertion (b) of the Corollary is equivalent to the existence of a path in $\underline{\text{KLND}}(B)$ going from A_1 to A_2 . Paragraph [5, 3.2.2] also defines a subgraph $\underline{\text{KLND}}_R(B)$ of the graph $\underline{\text{KLND}}(B)$, and clearly A_1, A_2 are two vertices of $\underline{\text{KLND}}_R(B)$; so, to prove (b), it suffices to show that $\underline{\text{KLND}}_R(B)$ is a connected graph. We have $R \in \mathcal{R}^{\text{in}}(B)$ (cf. [5, 5.2]) and consequently (cf. [5, 5.3], using that B is a UFD) we have an isomorphism of graphs $\underline{\text{KLND}}_R(B) \cong \underline{\text{KLND}}_K(\mathcal{B})$. As $\mathcal{B} \in \mathfrak{D}(K)$ by part (a), we may apply [5, 4.8] and conclude that $\underline{\text{KLND}}_K(\mathcal{B})$ is connected. Assertion (b) is proved. \square

The following is a trivial consequence of Corollary 4.4.

4.5. Corollary. *Let $B \in \mathcal{N}(\mathbf{k})$, where \mathbf{k} is a field of characteristic zero. Suppose that B has transcendence degree two over $\text{ML}(B)$.*

- (1) *Let $R = \text{ML}(B)$ and $K = \text{Frac } R$. Then $K \otimes_R B \in \mathfrak{D}(K)$.*
- (2) *If B is a UFD then, for any $A_1, A_2 \in \text{KLND}(B)$, there exists a finite sequence of local slice constructions which transforms A_1 into A_2 .*

REFERENCES

1. T. Bandman and L. Makar-Limanov, *Affine surfaces with $AK(S) = \mathbb{C}$* , Michigan Math. J. **49**, 2001, 567–582
2. J. Bertin, *Pinceaux de droites et automorphismes des surfaces affines*, J. Reine Angew. Math. **341** (1983), 32–53.
3. N. Bourbaki, *Éléments de mathématique. Algèbre. Chapitres 1 à 3*, Hermann, Paris, 1970.
4. D. Daigle, *On locally nilpotent derivations of $k[X_1, X_2, Y]/(\varphi(Y) - X_1X_2)$* , J. Pure and Appl. Algebra **181** (2003), 181–208.
5. D. Daigle, *Locally nilpotent derivations and Danielewski surfaces*, Osaka J. Math. **41** (2004), 37–80.
6. D. Daigle, *On polynomials in three variables annihilated by two locally nilpotent derivations*, J. of Algebra **310** (2007), 303–324.
7. D. Daigle and P. Russell, *On log \mathbb{Q} -homology planes and weighted projective planes*, Canadian J. of Math. **56** (2004), 1145–1189.
8. A. Dubouloz, *Completions of normal affine surfaces with a trivial Makar-Limanov invariant*, Michigan Math. J. **52** (2004), 289–308.
9. A. Dubouloz, *Danielewski-fieseler surfaces*, Transformation Groups **10** (2005), 139–162.
10. A. Dubouloz, *Embeddings of Danielewski surfaces in affine spaces*, Comment. Math. Helv. **81** (2006), 49–73.
11. A. van den Essen, *Polynomial automorphisms*, Progress in Mathematics, vol. 190, Birkhäuser, 2000.
12. G. Freudenburg, *Local slice constructions in $K[X, Y, Z]$* , Osaka J. Math. **34** (1997), 757–767.
13. G. Freudenburg, *Algebraic Theory of Locally Nilpotent Derivations*, Invariant theory and algebraic transformation groups VII, Springer-Verlag, 2006.
14. R.V. Gurjar, M. Miyanishi, *Automorphisms of affine surfaces with \mathbb{A}^1 -fibrations*, Michigan Math. J. **53** (2005), 33–55.
15. R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics, No. 52, Springer-Verlag, 1977.

16. L. Makar-Limanov, *On groups of automorphisms of a class of surfaces*, Israel J. of Math. **69** (1990), 250–256.
17. K. Masuda and M. Miyanishi, *The additive group actions on \mathbb{Q} -homology planes*, Ann. Inst. Fourier (Grenoble) **53** (2003), 429–464.
18. H. Matsumura, *Commutative Algebra*, 2nd edition, Benjamin Cummings, 1980, Mathematics Lecture Note Series.
19. Y. Nouazé and P. Gabriel, *Idéaux premiers de l'algèbre enveloppante d'une algèbre de Lie nilpotente*, J. of Algebra **6** (1967), 77–99.
20. R. Rentschler, *Opérations du groupe additif sur le plan affine*, C. R. Acad. Sc. Paris **267** (1968), 384–387.
21. D. Wright, *On the jacobian conjecture*, Illinois J. of Math. **25** (1981), 423–440.
22. O. Zariski and P. Samuel, *Commutative algebra, vol. 1*, Graduate Texts in Mathematics, No. 28, Springer-Verlag, New York, 1975.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA, CANADA
K1N 6N5

E-mail address: ddaigle@uottawa.ca